

Gaugings at angles from orientifold reductions

Diederik Roest

*Centre for Theoretical Physics,
University of Groningen,
Nijenborgh 4, 9747 AG Groningen, The Netherlands
d.roest@rug.nl*

ABSTRACT

We consider orientifold reductions to $\mathcal{N} = 4$ gauged supergravity in four dimensions. A special feature of this theory is that different factors of the gauge group can have relative angles with respect to the electro-magnetic $SL(2)$ symmetry. These are crucial for moduli stabilisation and De Sitter vacua. We show how such gaugings at angles generically arise in orientifold reductions.

1 Introduction

An important issue in string theory is the stabilisation of moduli. Compactifications to four dimensions generally lead to an abundance of scalar fields, which need to be stabilised at some point in moduli space. Flux compactifications are an attractive route to such a scenario [1]. In addition one would like to accomodate for a positive value of the scalar potential in this vacuum. Although at first this seemed hard to realise, there are now a number of possible models for De Sitter space-times within string theory [2].

Parallel to the ‘top-down’ approach of string compactifications one can also take a ‘bottom-up’ perspective. There have been systematic investigations of the possibilities for moduli stabilisation and De Sitter vacua in four-dimensional gauged supergravity, irrespective of any higher-dimensional origin. For $\mathcal{N} \geq 4$ extended supergravity, the De Sitter vacua found so far are unstable and have a value for the slow-roll parameter of order one [3, 4]. For $\mathcal{N} = 2$, on the other hand, there are examples with stable De Sitter vacua [5]. The higher-dimensional origin and relation to string theory of these cases is unknown.

In this paper we focus on $\mathcal{N} = 4$ supergravity, since the relevant aspects are very clear in that case. Both moduli stabilisation and De Sitter vacua crucially depend on a specific property of the gauging. First of all, the gauge group needs to be a product of factors. In addition, these gauge factors need to have different angles with respect to the electro-magnetic $SL(2)$ symmetry that rotates vectors into their electro-magnetic duals [6]. As will be discussed in more detail later, without such a structure the scalar potential V has an overall exponential dependence on the dilaton, making it impossible to stabilise the dilaton at a finite value of V . Therefore it is crucial to have a product of gauge factors with relative $SL(2)$ angles, i.e. *gaugings at angles*.

Despite many results on the relation between $\mathcal{N} = 4$ gaugings and their higher-dimensional ancestors, see e.g. [7–10], the higher-dimensional origin of non-trivial $SL(2)$ angles has never been clearly pointed out¹. In this paper we work out in detail a simple orientifold reduction and identify the resulting $\mathcal{N} = 4$ supergravity. The latter turns out to have gaugings at angles, thus providing a higher-dimensional origin for this feature. In particular, we show how moduli stabilisation is achieved by combining contributions to the scalar potential that originate from the bulk action and from the local source terms due to the orientifold. By clarifying the relation between gaugings at angles and orientifold reductions we aim to close the gap between the ‘top-down’

¹It was anticipated in [11] that orientifold reductions involving the Romans’ mass parameter and NS-NS flux would lead to non-trivial $SL(2)$ angles. However, no orientifold contributions and tadpole conditions were included (this was done subsequently for $\mathcal{N} = 1$ in [12]). More recently, the connection between orientifold reductions of massive IIA and non-trivial $SL(2)$ angles was conjectured in [13].

and ‘bottom-up’ approaches.

The organisation of this paper is as follows. In section 2, we review a number of general aspects of $\mathcal{N} = 4$ supergravity, after which we focus on a particular(ly useful) truncation. The structure of the gauging and scalar potential is emphasised. Section 3 discusses the orientifold reduction of IIA. Again we restrict ourselves to the simple truncation and show the equivalence to a specific $\mathcal{N} = 4$ theory. Finally, section 4 contains our conclusions and a number of remarks on possible extensions and the relation to other work.

2 $\mathcal{N} = 4$ gauged supergravity

In this section we discuss the structure of the $\mathcal{N} = 4$ theory and its gaugings. We will briefly summarise the general case and focus on a simple truncation which, while technically almost trivial, nevertheless retains the special feature of gaugings at angles that we want to highlight. In the next section this will be related to a simple orientifold reduction of IIA. For the general $\mathcal{N} = 4$ discussion we follow the conventions of [9], where more details and further references can be found.

The scalars of $D = 4$, $\mathcal{N} = 4$ supergravity parametrise a scalar coset of the form

$$\frac{SL(2)}{SO(2)} \times \frac{SO(6, 6+n)}{SO(6) \times SO(6+n)}, \quad (2.1)$$

The first factor contains the scalars of the supergravity multiplet. It is denoted by $M_{\alpha\beta}$, for which we use the following explicit parametrisation:

$$M_{\alpha\beta} = e^\phi \begin{pmatrix} \chi^2 + e^{-2\phi} & -\chi \\ -\chi & 1 \end{pmatrix}, \quad \alpha = (+, -). \quad (2.2)$$

The $SL(2)$ indices are raised and lowered with $\epsilon_{\alpha\beta} = \epsilon^{\alpha\beta}$, where $\epsilon^{+-} = -\epsilon^{-+} = 1$. The second factor in (2.1) is spanned by the matter multiplets. We focus on the case of six matter multiplets, corresponding to $n = 0$. In this case it is convenient to use light-cone coordinates for the $SO(6, 6)$ group. The invariant metric is of the form

$$\eta_{MN} = \eta^{MN} = \begin{pmatrix} & \mathbb{I}_6 \\ \mathbb{I}_6 & \end{pmatrix}, \quad M = (1, \dots, 6, \bar{1}, \dots, \bar{6}). \quad (2.3)$$

The corresponding $SO(6, 6)$ element that parametrises the scalar coset is denoted by M_{MN} . We will introduce an explicit parametrisation later. Together with the Einstein-Hilbert term for the metric, the scalars have the following kinetic terms²:

$$\mathcal{L}_{\text{kin}} = \sqrt{-g} [R + \tfrac{1}{4} \partial_\mu M_{\alpha\beta} \partial^\mu M^{\alpha\beta} + \tfrac{1}{8} \partial_\mu M_{MN} \partial^\mu M^{MN}]. \quad (2.4)$$

²We have multiplied the total action of [9] with a factor of two.

In addition the theory contains $12 + n$ vectors, transforming in the fundamental representation of $SO(6, 6 + n)$. A noteworthy feature is that under the compact part of the $SL(2)$ symmetry these transform into their electro-magnetic dual. This symmetry is therefore only realised on-shell. This is a particular feature of four-dimensional theories and leads to the following intricate structure of gaugings.

The possible gaugings of this theory have been classified within the framework of the embedding tensor [14]. It turns out that one can introduce two $SO(6, 6)$ representations of gauge parameters: an anti-symmetric three-form $f_{\alpha MNP}$ and a fundamental $\xi_{\alpha M}$, both of which transform as a doublet under $SL(2)$. Consistency of such gaugings requires a number of quadratic constraints on the embedding tensor, which can be seen as generalised Jacobi identities. For later purposes we will give the constraints for the case with $\xi_{\alpha M} = 0$, for which one finds

$$f_{\alpha R[MN}f_{\beta PQ]}{}^R = 0, \quad \epsilon^{\alpha\beta}f_{\alpha MNR}f_{\beta PQ}{}^R = 0. \quad (2.5)$$

The combination of supersymmetry and gaugings induce the following scalar potential:

$$\begin{aligned} \mathcal{L}_{\text{pot}} &= -\sqrt{-g}V, \\ V &= \frac{1}{8}f_{\alpha MNP}f_{\beta QRS}M^{\alpha\beta}[\frac{1}{3}M^{MQ}M^{NR}M^{PS} + (\frac{2}{3}\eta^{MQ} - M^{MQ})\eta^{NR}\eta^{PS}] + \\ &\quad - \frac{1}{18}f_{\alpha MNP}f_{\beta QRS}\epsilon^{\alpha\beta}M^{MNPQRS} + \frac{3}{8}\xi_{\alpha}^M\xi_{\beta}^NM^{\alpha\beta}M_{MN}, \end{aligned} \quad (2.6)$$

where the definition of M^{MNPQRS} in terms of M^{MN} can be found in [9].

As mentioned before, an important aspect of this four-dimensional supergravity is that vectors are transformed into their electro-magnetic dual under the $SO(2) \subset SL(2)$ symmetry. This on-shell symmetry is responsible for the $SL(2)$ doublet structure of the gauge parameters. Depending on the $SL(2)$ orientation, the embedding tensor picks out a vector or its dual (or a linear combination) to gauge a part of the global symmetry of the theory. Moreover, when the gauge group is a product of different factors, it is possible to choose a different $SL(2)$ orientation for the different factors. In terms of the embedding tensor, this corresponds to

$$f_{\alpha MNP} = \sum_i \delta_{\alpha}^{(i)} f_{MNP}^{(i)}, \quad \xi_{\alpha M} = \sum_i \delta_{\alpha}^{(i)} \xi_M^{(i)}, \quad (2.7)$$

where $f_{MNP}^{(i)}$ and $\xi_M^{(i)}$ specify a factor of the gauge group and the $\delta_{\alpha}^{(i)}$ do not necessarily point in the same $SL(2)$ directions. This possibility is referred to as different $SL(2)$ (or $SU(1, 1)$) or De Roo-Wagemans angles [6]. If all the $SL(2)$ factors are identical, one can always rotate these to the $\alpha = +$ direction, corresponding to a zero angle. It follows from (2.6) that in such cases the scalar potential has an overall dependence of e^{ϕ} and hence a runaway direction. Therefore gaugings at angles play a crucial role in moduli stabilisation [4].

Instead of the full $\mathcal{N} = 4$ supergravity we consider the following truncation. The $SO(6, 6)$ symmetry can be decomposed into

$$SL(3) \times SL(3) \times \mathbb{R}^+ \times \mathbb{R}^+ \subset SL(6) \times \mathbb{R}^+ \subset SO(6, 6). \quad (2.8)$$

We focus on the subsector of the theory that is invariant under both $SL(3)$ factors. The group-theoretic nature of this truncation guarantees its consistency.

A drastic consequence is that all vectors are projected out. This follows from the decomposition of the fundamental representation of $SO(6, 6)$ into $SL(3) \times SL(3)$ (omitting the $\mathbb{R}^+$ weights):

$$\mathbf{12} \rightarrow (\mathbf{3}, \mathbf{1}) \oplus (\mathbf{3}', \mathbf{1}) \oplus (\mathbf{1}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{3}'), \quad (2.9)$$

where no singlets appear.

In the scalar sector, the $SL(2)$ scalars are unaffected by this truncation. In contrast, from the decomposition of the adjoint representation one learns that many of the $SO(6, 6)$ scalars are projected out:

$$\mathbf{66} \rightarrow (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}', \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}') \oplus (\mathbf{1}, \mathbf{8}) \oplus (\mathbf{8}, \mathbf{1}). \quad (2.10)$$

Since there are only two singlets, the truncation preserves two dilatonic scalars. One can take the following parametrisation of the $SO(6, 6)$ element M_{MN} in terms of these scalars φ_1 and φ_2 :

$$M_{MN} = \begin{pmatrix} e^{-\sqrt{2/3} \varphi_1} & & & \\ & e^{-\sqrt{2/3} \varphi_2} & & \\ & & e^{\sqrt{2/3} \varphi_1} & \\ & & & e^{\sqrt{2/3} \varphi_2} \end{pmatrix} \otimes \mathbb{I}_3. \quad (2.11)$$

Inserting this in (2.4) gives rise to the following kinetic terms for the four scalars that survive the truncation:

$$\mathcal{L}_{\text{kin}} = \sqrt{-g} [R - \frac{1}{2}(\partial\phi)^2 - \frac{1}{2}e^{2\phi}(\partial\chi)^2 - \frac{1}{2}(\partial\varphi_1)^2 - \frac{1}{2}(\partial\varphi_2)^2], \quad (2.12)$$

where we have also included the Einstein-Hilbert term for the metric.

We now come to effect of the truncation of the embedding tensor. As the components $\xi_{\alpha M}$ also transform in the fundamental representation of $SO(6, 6)$, these suffer the same fate as the vectors, and are all projected out. The other components $f_{\alpha MNP}$ give rise to a number of $SL(3) \times SL(3)$ representations, including four singlets (omitting the $SL(2)$ doublet structure):

$$\mathbf{220} \rightarrow 4 \cdot (\mathbf{1}, \mathbf{1}) \oplus \text{non-singlet representations}, \quad (2.13)$$

So there are four $SO(6,6)$ components that survive the truncation. In our light-cone basis these correspond to $f_{\alpha 123}$, $f_{\alpha 456}$, $f_{\alpha \bar{1}\bar{2}\bar{3}}$ and $f_{\alpha \bar{4}\bar{5}\bar{6}}$. Moreover, the quadratic constraints (2.5) result in the simple conditions

$$f_{\alpha 123} f_{\beta \bar{1}\bar{2}\bar{3}} = 0, \quad f_{\alpha 456} f_{\beta \bar{4}\bar{5}\bar{6}} = 0. \quad (2.14)$$

Hence there are four possibilities of gauge parameters in this truncated theory, taking either non-zero ‘unbarred’ or ‘barred’ components in the (123) and independently in the (456) directions.

These four models can in fact be related to each other by particular elements of the global symmetry, which interchange the two types of light-cone directions. For example, an $SO(6,6)$ transformation of the form (2.3) interchanges the six ‘unbarred’ and ‘barred’ directions. For an odd number of interchanged directions this transformation needs to be accompanied by a sign flip of the $SL(2)$ axion. Therefore the four models are physically equivalent, and in the following we will only consider the case with gauge parameters $f_{\alpha 123}$ and $f_{\beta 456}$.

Note that the model has a product of gauge factors³: one in the (123) directions and one in the (456) directions. These gaugings are specified by four real parameters: two can be seen as gauge coupling constants while the other two correspond to the $SL(2)$ angles of the two gauge factors. One of the angles can be set equal to zero, i.e. point in the $\alpha = +$ direction, by an $SO(2) \subset SL(2)$ transformation. We will use this to rotate away f_{-123} . Moreover, if f_{-456} does not vanish, one can perform an $SL(2)$ transformation that shifts the axion to set the second angle to 90 degrees. This corresponds to setting $f_{+456} = 0$. In the case of two different angles, one can therefore always take these orthogonal. We will not use this and keep the second angle arbitrary, however.

Let us analyse the form of the scalar potential and its extrema for the truncated model. By writing out the general scalar potential (2.6) and using $f_{-123} = 0$, one finds:

$$V = \frac{1}{4} \left(f_{+123} e^{\phi/2 + \sqrt{3/2} \varphi_1} - f_{-456} e^{-\phi/2 + \sqrt{3/2} \varphi_2} \right)^2 + \frac{1}{4} \left(f_{+456} + \chi f_{-456} \right)^2 e^{\phi + \sqrt{6} \varphi_2}. \quad (2.15)$$

Strikingly, the potential combines into the sum of two squares and is positive definite. This relies crucially on the different $SL(2)$ angles: the crossterm in the first square is independent of the $SL(2)$ scalars and comes from the last line of (2.6). Only in the presence of such terms can one have moduli stabilisation. In the extremum with respect

³This is a slight abuse of notation, as the truncated model does not have any vectors. However, by including the fields that have been truncated out, this model can be restored to a unique $\mathcal{N} = 4$ supergravity with a gauging defined by these parameters.

to χ , the term on the second line vanishes. The remaining square has an extremum if and only if f_{+123} and f_{-456} have equal signs, while the dilatonic scalars are such that the first square in the potential vanishes as well. In the extremum both squares that make up the scalar potential vanish, and we have a Minkowski solution.

Next, we investigate the issue of stability. The axion can be seen to decouple from this issue as $\delta_\chi \delta_\chi V$ is positive while $\delta_\chi \delta_{\vec{\phi}} V$ vanishes, where $\vec{\phi}$ represents the three dilatonic scalars and $\delta_\chi \equiv \delta/\delta\chi$ etc. The matrix $\delta_{\vec{\phi}} \delta_{\vec{\phi}} V$ turns out to have one positive and two vanishing eigenvalues. The Minkowski solution is therefore a minimum of the scalar potential - at least in the truncation to $SL(3) \times SL(3)$ invariant scalars that we consider.

An interesting question is which gaugings are induced by gauge parameters of the form above. The answer can be found in [15], where so-called $CSO(p, q, r)$ -gaugings are considered. These groups can be seen as group contractions of $SO(p', q')$ with $p' + q' = p + q + r$. It turns out that each component of the structure constants that we consider induces a $CSO(1, 0, 3)$ gauging⁴ inside $SO(6, 6)$. Our gauge group therefore consists of a product of two such factors. The total dimension of these gauge groups is twelve, in accordance with the number of vectors. Reference [15] also performed a stability analysis with respect to all scalars and found a number of unstable directions. The Minkowski solution is therefore a saddlepoint of the full $\mathcal{N} = 4$ theory.

3 Orientifold reduction of IIA

In this section we will consider a simple orientifold reduction of the IIA theory, which will be related to the previous $\mathcal{N} = 4$ truncation. Further details on different aspects and more complicated cases can be found in e.g. [7, 8, 11, 12, 16].

Consider the toroidal reduction of massive IIA to four dimensions. Introducing an O6-plane corresponds to modding out by $(-)^{F_L} \Omega I_{7,8,9}$. Here $(-)^{F_L}$ and Ω correspond to the left-moving fermion number and world-sheet parity, respectively, whose combined action on the IIA bosonic fields is

$$\begin{aligned} \{\hat{g}_{\mu\nu}, \hat{\phi}, \hat{C}_3, \hat{C}_7\} &\rightarrow +\{\hat{g}_{\mu\nu}, \hat{\phi}, \hat{C}_3, \hat{C}_7\}, \\ \{\hat{B}, \hat{C}_1, \hat{C}_5, \hat{C}_9\} &\rightarrow -\{\hat{B}, \hat{C}_1, \hat{C}_5, \hat{C}_9\}. \end{aligned} \quad (3.1)$$

In addition, the space-time parity operation $I_{7,8,9}$ reverses the sign of three of the coordinates on the torus:

$$\{x^7, x^8, x^9\} \rightarrow -\{x^7, x^8, x^9\}. \quad (3.2)$$

⁴Modulo two typo's in these expressions, the $CSO(1, 0, 3)$ structure constants given in appendix B of [15] in a Cartesian basis correspond to f_{123} in our light-cone basis.

The indices in (3.1) are taken inside the O6-plane, i.e. in the directions $(0, \dots, 6)$. Other components with indices transverse to the O6-plane will acquire additional signs due to (3.2). Furthermore the Romans' mass parameter $\hat{G}_0$ of IIA is invariant under the above involution.

Instead of the general orientifold reduction we will focus on the following truncation. Consider the two T^3 's in the directions $\{x^4, x^5, x^6\}$ and $\{x^7, x^8, x^9\}$. Diffeomorphisms leaving the two factors separately invariant generate an $SL(3) \times SL(3)$ symmetry in the four-dimensional description. Completely analogous to the truncation of the $\mathcal{N} = 4$ theory of the previous section, we will retain only singlets with respect to both factors.

The most general Ansatz for the ten-dimensional metric that is consistent with $SL(3) \times SL(3)$ invariance is of the form

$$\hat{g}_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} e^{\sqrt{3}/2 \sigma_2} g_{\mu\nu} & & \\ & e^{-\sigma_2/2\sqrt{3} + \sigma_3/\sqrt{3}} \mathbb{I}_3 & \\ & & e^{-\sigma_2/2\sqrt{3} - \sigma_3/\sqrt{3}} \mathbb{I}_3 \end{pmatrix}, \quad (3.3)$$

consisting of the four-dimensional metric $g_{\mu\nu}$ and two scalars σ_2 and σ_3 . Both the Kaluza-Klein vectors and other scalars, parametrising deformations of the internal torus, are projected out by the truncation to $SL(3) \times SL(3)$ invariant fields. Furthermore, the normalisation of the σ 's is chosen to ensure canonical normalisation. The ten-dimensional bulk action is⁵

$$\hat{S} = 2\pi \int d^{10}x (\hat{\mathcal{L}}_1 + \hat{\mathcal{L}}_2), \quad (3.4)$$

where the first term contains the Einstein-Hilbert term and the dilaton kinetic term, while the second term is concerned with the gauge potentials. For the first part, after reduction to four dimensions we find

$$\hat{\mathcal{L}}_1 = \sqrt{-\hat{g}} [\hat{R} - \frac{1}{2}(\partial\hat{\phi})^2] \quad \rightarrow \quad \mathcal{L}_1 = \sqrt{-g} [R - \frac{1}{2} \sum_{i=1,2,3} (\partial\sigma_i)^2]. \quad (3.5)$$

where we have set $\hat{\phi} = \sigma_1$. Note that we use Einstein frame both in ten and in four dimensions.

Next, we turn to the gauge potentials. The NS-NS two-form potential is odd under (3.1) and hence has to wrap an odd cycle in the torus. However, there is only one such form that is invariant under $SL(3) \times SL(3)$: a three-form. The field strength of this gauge potential therefore only gives a constant parameter h_3 :

$$\hat{H} = d\hat{B} = h_3 dx^7 \wedge dx^8 \wedge dx^9. \quad (3.6)$$

⁵Our IIA conventions agree with e.g. [17]. To avoid cluttering our formulae we have set $4\pi^2\alpha' = 1$.

The R-R gauge potentials are either even or odd. First of all, the Romans' mass parameter, which can be seen as a zero-form R-R field strength, is even and also gives rise to a constant parameter in four dimensions: $\hat{G}_0 = g_0$. The R-R vector is odd under the orientifold involution. Its field strength necessarily vanishes,

$$\hat{G}_2 = d\hat{C}_1 + \hat{G}_0 \hat{B} = 0, \quad (3.7)$$

as there are no odd $SL(3) \times SL(3)$ invariant zero-, one- or two-cycles on the torus. Finally, the R-R three-form is even. Its magnetic part will be proportional to the even $SL(3) \times SL(3)$ invariant three-form and give rise to a scalar χ ,

$$\hat{G}_4^{(m)} = d\hat{C}_3 - \hat{H} \wedge \hat{C}_1 + \frac{1}{2} \hat{B} \wedge B = d\chi \wedge dx^4 \wedge dx^5 \wedge dx^6. \quad (3.8)$$

It can also have an electric part. This will be more conveniently described in terms of the dual field strength, which is related by $\hat{G}_6^{(m)} = e^{\hat{\phi}/2} \star \hat{G}_4^{(e)}$. The dual five-form gauge potential is odd under (3.1) and can wrap the total six-torus:

$$\hat{G}_6^{(m)} = d\hat{C}_5 - \hat{H} \wedge \hat{C}_3 + \frac{1}{6} \hat{B} \wedge B \wedge B = (g_6 + h_3 \chi) dx^4 \wedge \cdots \wedge dx^9. \quad (3.9)$$

Quantisation of these parameters requires g_0 , h_3 and g_6 all to be integer. Moreover, we will assume $g_0 h_3$ to be positive, for reasons that will become clear later.

With the Ansätze above, the kinetic terms for the IIA gauge potentials reduce to a kinetic term for χ and potential terms for the three constants h_3 , g_0 and g_6 :

$$\begin{aligned} \hat{\mathcal{L}}_2 &= \sqrt{-\hat{g}} \left[-\frac{1}{2} e^{-\hat{\phi}} \hat{H} \cdot \hat{H} - \frac{1}{2} e^{5/2 \hat{\phi}} \hat{G}_0^2 - \frac{1}{2} e^{\hat{\phi}/2} \hat{G}_4^{(m)} \cdot \hat{G}_4^{(m)} - \frac{1}{2} e^{-\hat{\phi}/2} \hat{G}_6^{(m)} \cdot \hat{G}_6^{(m)} \right] \rightarrow \\ \mathcal{L}_2 &= \sqrt{-g} \left[-\frac{1}{2} e^{\sigma_1/2 + \sqrt{3}/2 \sigma_2 - \sqrt{3} \sigma_3} (\partial\chi)^2 - \frac{1}{2} h_3^2 e^{-\sigma_1 + \sqrt{3}(\sigma_2 + \sigma_3)} - \frac{1}{2} g_0^2 e^{5/2 \sigma_1 + \sqrt{3}/2 \sigma_2} + \right. \\ &\quad \left. - \frac{1}{2} (g_6 + h_3 \chi)^2 e^{-\sigma_1/2 + 3\sqrt{3}/2 \sigma_2} \right]. \end{aligned} \quad (3.10)$$

Note that there are no topological Chern-Simons terms in the democratic formulation of IIA [18]; the kinetic terms for the different R-R potentials suffice. These are therefore all the contributions from the ten-dimensional bulk action.

In addition to the bulk, one must also include the orientifold planes induced by (3.1) and (3.2). We further allow for a number of D6-branes with the same orientation (ignoring the world-volume excitations). These give rise to the following contributions to the scalar potential:

$$\hat{S}_{\text{O6/D6}} = 2\pi N \int d^7x \left[e^{3/4 \hat{\phi}} \sqrt{-\hat{g}_7} \right] \rightarrow \mathcal{L}_3 = \sqrt{-g} \left[N e^{3/4 \sigma_1 + 3\sqrt{3}/4 \sigma_2 + \sqrt{3}/2 \sigma_3} \right], \quad (3.11)$$

where $N = 2N_{\text{O6}} - N_{\text{D6}}$. An orientifolded three-torus has 2^3 fixed points under (3.2) and would lead to $N_{\text{O6}} = 8$. The unusual dilaton coupling stems from the fact that we

are using Einstein frame. Furthermore, we have not included the Wess-Zumino term, as this will not contribute to the four-dimensional action. The total resulting action consists of the three pieces (3.5), (3.10) and (3.11).

The Bianchi identities for the different field strengths read

$$d\hat{H} = 0, \quad d\hat{G}_{2n+2} = \hat{H} \wedge \hat{G}_{2n}. \quad (3.12)$$

These are satisfied by the Ansätze above modulo the following two points. The first is that, due to the Wess-Zumino term of the O6-planes and D6-branes that involves $\hat{C}_7$, the Bianchi identity of $\hat{G}_2$ is modified:

$$d\hat{G}_2 = \hat{G}_0 \hat{H} - N dx^7 \wedge dx^8 \wedge dx^9, \quad \Rightarrow \quad g_0 h_3 = N, \quad (3.13)$$

leading to a tadpole condition that will be essential. Furthermore, the reader might worry about the Bianchi identity for the electric part of $\hat{G}_6$, which does not vanish identically. However, this will be proportional to the four-dimensional field equation for χ and vanishes on-shell.

Turning to the three pieces of which the action consists, we can now appreciate the beauty of the orientifold reduction and the underlying supersymmetry. The contribution due to the orientifold is such that the scalar potential terms (3.10) and (3.11) involving g_0 and h_3 can be combined into a square. This crucially relies on the tadpole condition (3.13). The scalar potential is now a positive definite sum of two squares.

The orientifold breaks half of supersymmetry and the resulting four-dimensional description is an $\mathcal{N} = 4$ supergravity. Since our truncation to $SL(3) \times SL(3)$ singlets coincides with that of the previous section, there must be a relation to the model discussed there. Indeed the two can be related by the following field redefinition for the σ 's:

$$\begin{pmatrix} \phi \\ \varphi_1 \\ \varphi_2 \end{pmatrix} = \frac{1}{4\sqrt{2}} \begin{pmatrix} \sqrt{2} & 3\sqrt{3} & -\sqrt{3} \\ \sqrt{6} & 1 & 5 \\ -2\sqrt{6} & 2 & 2 \end{pmatrix} \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix}, \quad (3.14)$$

in terms of the $SL(2)$ dilaton ϕ and the $SO(6,6)$ dilatons φ_1 and φ_2 of the previous section. Furthermore, one must identify the gauge parameters of both models as

$$(f_{+123}, f_{-456}, f_{+456}) = \sqrt{2}(g_0, h_3, g_6). \quad (3.15)$$

These redefinitions turn the Lagrangian consisting of (2.12) and (2.15) into that consisting of (3.5), (3.10) and (3.11). Moreover, since the $SL(3) \times SL(3)$ invariant model defines a unique $\mathcal{N} = 4$ gauged supergravity, this connection extends to the full theory: an orientifold reduction that retains all fields and includes these three fluxes will lead to an $\mathcal{N} = 4$ supergravity with gauge parameters (3.15).

Our simple orientifold reduction therefore leads to an $\mathcal{N} = 4$ supergravity with $CSO(1, 0, 3) \times CSO(1, 0, 3)$ gauge group, where the two gauge factors have a non-vanishing relative $SL(2)$ angle. Note that the tadpole condition implies a relation on the gauge parameters: they have to be of the form (3.15) with g_0, h_3, g_6 integer and subject to $g_0 h_3 = N$. Furthermore, the condition on the signs of f_{+123} and f_{-456} of the previous section justifies our assumption that $g_0 h_3$ is positive.

4 Discussion and outlook

In the previous sections we have seen that the simple IIA orientifold reduction with fluxes (g_0, h_3, g_6) leads to the $\mathcal{N} = 4$ supergravity with $CSO(1, 0, 3) \times CSO(1, 0, 3)$ gauge group of [15]. The two gauge factors generically are at a non-vanishing $SL(2)$ angle with respect to each other, leading to moduli stabilisation. From the orientifold side, this important feature was achieved by a collaboration of contributions to the scalar potential from the IIA bulk action (3.10) and the local source terms (3.11) due to the orientifolding. In order to avoid this, one must tune the O6/D6 content such that $N = 0$, in which case the two gauge groups have the same angle or one of them disappears. Thus we have clarified the higher-dimensional origin of the important $\mathcal{N} = 4$ phenomenon of $SL(2)$ angles. Our simple model demonstrates that such gaugings at angles will be a generic outcome of IIA orientifold reductions.

Due to T-duality our results can be related to other orientifold cases. For instance, consider the case where we T-dualise in the three toroidal directions (x^4, x^5, x^6) of the O6-plane worldvolume. The resulting IIB reduction involves an O3-plane and has been studied at length in e.g. [7, 19]. Our results have a clear counterpart in this IIB case. The parameters g_0, h_3 and g_6 now come from the IIB three-form components $\hat{G}_{456}, \hat{H}_{789}$ and $\hat{G}_{789}$, respectively. The tadpole condition relates D3-branes and O3-planes to a contribution due to the complex three-form flux, and the resulting action also contains a sum of squares. In this case the vanishing of the squares corresponds to the well-known imaginary self-duality condition on the three-form flux. Again the non-trivial $SL(2)$ angles play an important role in the stabilisation of moduli.

On the other hand, one could consider T-duality in any of the directions (x^7, x^8, x^9) transverse to the O6-plane. In contrast to the previous case, T-duality in these directions does not leave the three-form flux invariant. Instead it has been argued that this will be transformed into geometric or even non-geometric flux [20]. Therefore T-duality in the transverse directions, giving rise O7-, O8- or O9-planes, does not lead to the simple reductions we considered with only gauge fluxes.

Coming back to the O6-plane, the reduction to four dimensions can in fact be split

up in two steps. The first consists of the reduction over the transverse space of the orientifold, while the second reduces over its three toroidal world-volume directions. One could stop after the first step and thus obtain a seven-dimensional half-maximal supergravity theory with two parameters g_0 and h_3 (the remaining g_6 only shows up after the second step). The gaugings of this theory are encoded in representations ξ_m and f_{mnp} of $SO(3,3)$, while there is a single topological mass parameter m [21]. Restricting to $SL(3, \mathbb{R})$ invariant components leads to f_{123} and $f_{\bar{1}\bar{2}\bar{3}}$ in addition to the mass parameter. The product of the two gauge parameters vanishes due to the Jacobi identity. Therefore, the orientifold parameters g_0 and h_3 are to be identified with m and e.g. f_{123} . A subsequent normal toroidal reduction to four dimensions leads to the theory that we have considered in this paper.

Two lessons can be drawn from this discussion. Firstly, the $D = 7$ topological mass parameter m has a higher-dimensional origin from orientifold reductions. Secondly, gaugings at angles in four dimensions are induced by a toroidal reduction of the seven-dimensional massive theory. To our knowledge, this is the first time that a higher-dimensional origin of gaugings at angles from $4 < D \leq 7$ half-maximal supergravity has been put forward. It would be interesting to investigate this connection in more detail. Due to the above discussion involving Op -planes with $p > 6$ we do not expect such an origin from dimensions higher than seven. This ties in nicely with the absence of mass parameters in these theories [21].

In this paper we have restricted ourselves to a very simple truncation to $SL(3) \times SL(3)$ singlets. Needless to say this can be relaxed to allow for many more possibilities [7, 8]: different components of gauge fluxes can be turned on and one could reduce over twisted tori with non-vanishing geometric fluxes ω . This would lead to additional structure constants, inducing different gaugings of the four-dimensional $\mathcal{N} = 4$ theory. For instance, including ω and $\hat{G}_2$ fluxes in a specific way could lead to $CSO(3, 0, 1) \times CSO(3, 0, 1)$ gaugings [11]. It would be interesting to investigate a possible relation to the $SU(2) \times SU(2)$ reduction of [22]. Furthermore, such reductions might give a higher-dimensional origin to the unstable De Sitter vacua of [4].

Finally, one can consider orientifold reductions that break more supersymmetry. It would be of great interest if one could find e.g. a string-theoretic origin for the stable De Sitter vacua in $\mathcal{N} = 2$ supergravity [5], for which non-trivial $SL(2)$ angles are a necessary ingredient.

Acknowledgements

We thank Giuseppe Dibitetto, Thomas van Riet and Mees de Roo for very stimulating discussions. The work of DR is supported by a VIDI grant from the Netherlands Organisation for Scientific Research (NWO).

References

- [1] This has been reviewed in e.g.
M. Grana, *Flux compactifications in string theory: A comprehensive review*, Phys. Rept. **423** (2006) 91–158 [[hep-th/0509003](#)].
M. R. Douglas and S. Kachru, *Flux compactification*, Rev. Mod. Phys. **79** (2007) 733–796 [[hep-th/0610102](#)].
- [2] Early references are e.g.
S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, *De Sitter vacua in string theory*, Phys. Rev. **D68** (2003) 046005 [[hep-th/0301240](#)].
C. P. Burgess, R. Kallosh and F. Quevedo, *de Sitter String Vacua from Supersymmetric D-terms*, JHEP **10** (2003) 056 [[hep-th/0309187](#)].
- [3] R. Kallosh, A. D. Linde, S. Prokushkin and M. Shmakova, *Gauged supergravities, de Sitter space and cosmology*, Phys. Rev. **D65** (2002) 105016 [[hep-th/0110089](#)].
- [4] M. de Roo, D. B. Westra and S. Panda, *De Sitter solutions in $N = 4$ matter coupled supergravity*, JHEP **02** (2003) 003 [[hep-th/0212216](#)].
M. de Roo, D. B. Westra, S. Panda and M. Trigiante, *Potential and mass-matrix in gauged $N = 4$ supergravity*, JHEP **11** (2003) 022 [[hep-th/0310187](#)].
- [5] P. Fre, M. Trigiante and A. Van Proeyen, *Stable de Sitter vacua from $N = 2$ supergravity*, Class. Quant. Grav. **19** (2002) 4167–4194 [[hep-th/0205119](#)].
O. Ogetbil, *Stable de Sitter Vacua in 4 Dimensional Supergravity Originating from 5 Dimensions*, Phys. Rev. **D78** (2008) 105001 [[0809.0544](#)].
- [6] M. de Roo and P. Wagemans, *Gauge matter coupling in $N=4$ supergravity*, Nucl. Phys. **B262** (1985) 644.
- [7] R. D’Auria, S. Ferrara and S. Vaula, *$N = 4$ gauged supergravity and a IIB orientifold with fluxes*, New J. Phys. **4** (2002) 71 [[hep-th/0206241](#)].

- R. D'Auria, S. Ferrara, F. Gargiulo, M. Trigiante and S. Vaula, *$N = 4$ supergravity Lagrangian for type IIB on $T^{**}6/Z(2)$ in presence of fluxes and D3-branes*, JHEP **06** (2003) 045 [[hep-th/0303049](#)].
- [8] C. Angelantonj, S. Ferrara and M. Trigiante, *New $D = 4$ gauged supergravities from $N = 4$ orientifolds with fluxes*, JHEP **10** (2003) 015 [[hep-th/0306185](#)].
C. Angelantonj, S. Ferrara and M. Trigiante, *Unusual gauged supergravities from type IIA and type IIB orientifolds*, Phys. Lett. **B582** (2004) 263–269 [[hep-th/0310136](#)].
G. Dall'Agata and S. Ferrara, *Gauged supergravity algebras from twisted tori compactifications with fluxes*, Nucl. Phys. **B717** (2005) 223–245 [[hep-th/0502066](#)].
- [9] J. Schon and M. Weidner, *Gauged $N = 4$ supergravities*, JHEP **05** (2006) 034 [[hep-th/0602024](#)].
- [10] J.-P. Derendinger, P. M. Petropoulos and N. Prezas, *Axionic symmetry gaugings in $N = 4$ supergravities and their higher-dimensional origin*, Nucl. Phys. **B785** (2007) 115–134 [[0705.0008](#)].
- [11] J.-P. Derendinger, C. Kounnas, P. M. Petropoulos and F. Zwirner, *Superpotentials in IIA compactifications with general fluxes*, Nucl. Phys. **B715** (2005) 211–233 [[hep-th/0411276](#)].
- [12] G. Villadoro and F. Zwirner, *$N = 1$ effective potential from dual type-IIA D6/O6 orientifolds with general fluxes*, JHEP **06** (2005) 047 [[hep-th/0503169](#)].
- [13] S. S. Haque, G. Shiu, B. Underwood and T. Van Riet, *Minimal simple de Sitter solutions*, Phys. Rev. D **79** (2009) 086005 [[0810.5328](#)].
- [14] H. Nicolai and H. Samtleben, *Compact and noncompact gauged maximal supergravities in three dimensions*, JHEP **04** (2001) 022 [[hep-th/0103032](#)].
B. de Wit, H. Samtleben and M. Trigiante, *On Lagrangians and gaugings of maximal supergravities*, Nucl. Phys. **B655** (2003) 93–126 [[hep-th/0212239](#)].
- [15] M. de Roo, D. B. Westra and S. Panda, *Gauging CSO groups in $N = 4$ supergravity*, JHEP **09** (2006) 011 [[hep-th/0606282](#)].
- [16] O. DeWolfe, A. Giryavets, S. Kachru and W. Taylor, *Type IIA moduli stabilization*, JHEP **07** (2005) 066 [[hep-th/0505160](#)].

- P. G. Camara, A. Font and L. E. Ibanez, *Fluxes, moduli fixing and MSSM-like vacua in a simple IIA orientifold*, JHEP **09** (2005) 013 [[hep-th/0506066](#)].
- [17] J. Polchinski, *String theory. Vol. 2: Superstring theory and beyond*, Cambridge, UK: Univ. Pr. (1998) 531 p.
- [18] E. Bergshoeff, R. Kallosh, T. Ortin, D. Roest and A. Van Proeyen, *New Formulations of D=10 Supersymmetry and D8-O8 Domain Walls*, Class. Quant. Grav. **18** (2001) 3359–3382 [[hep-th/0103233](#)].
- [19] S. Kachru, M. B. Schulz and S. Trivedi, *Moduli stabilization from fluxes in a simple IIB orientifold*, JHEP **10** (2003) 007 [[hep-th/0201028](#)].
A. R. Frey and J. Polchinski, *$N = 3$ warped compactifications*, Phys. Rev. **D65** (2002) 126009 [[hep-th/0201029](#)].
- [20] J. Shelton, W. Taylor and B. Wecht, *Nongeometric Flux Compactifications*, JHEP **10** (2005) 085 [[hep-th/0508133](#)].
- [21] E. A. Bergshoeff, J. Gomis, T. A. Nutma and D. Roest, *Kac-Moody Spectrum of (Half-)Maximal Supergravities*, JHEP **02** (2008) 069 [[0711.2035](#)].
- [22] C. Caviezel *et al.*, *On the Cosmology of Type IIA Compactifications on $SU(3)$ -structure Manifolds*, JHEP **04** (2009) 010 [[0812.3551](#)].